Global Change Biology

Global Change Biology (2016), doi: 10.1111/gcb.13299

Land-use change outweighs projected effects of changing rainfall on tree cover in sub-Saharan Africa

JULIE C. ALEMAN¹, OLIVIER BLARQUEZ² and CARLA A. STAVER¹ ¹Ecology and Evolutionary Biology, Yale University, P.O. Box 208106, New Haven, CT 06520, USA, ²Département de Géographie, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, QC H3C 3J7, Canada

Abstract

Global change will likely affect savanna and forest structure and distributions, with implications for diversity within both biomes. Few studies have examined the impacts of both expected precipitation and land use changes on vegetation structure in the future, despite their likely severity. Here, we modeled tree cover in sub-Saharan Africa, as a proxy for vegetation structure and land cover change, using climatic, edaphic, and anthropic data ($R^2 = 0.97$). Projected tree cover for the year 2070, simulated using scenarios that include climate and land use projections, generally decreased, both in forest and savanna, although the directionality of changes varied locally. The main driver of tree cover changes was land use change; the effects of precipitation change were minor by comparison. Interestingly, carbon emissions mitigation via increasing biofuels production resulted in decreases in tree cover, more severe than scenarios with more intense precipitation change, especially within savannas. Evaluation of tree cover change against protected area extent at the WWF Ecoregion scale suggested areas of high biodiversity and ecosystem services concern. Those forests most vulnerable to large decreases in tree cover were also highly protected, potentially buffering the effects of global change. Meanwhile, savannas, especially where they immediately bordered forests (*e.g.* West and Central Africa), were characterized by a dearth of protected areas, making them highly vulnerable. Savanna must become an explicit policy priority in the face of climate and land use change if conservation and livelihoods are to remain viable into the next century.

Keywords: habitat loss, land use, protected areas, rainfall, tree cover

Received 15 December 2015; revised version received 17 March 2016 and accepted 18 March 2016

Introduction

The tropical biosphere will potentially respond substantially to both climate and land use change. Existing predictions have focused mostly on the impacts of changing climate (Higgins & Scheiter, 2012; Zeng et al., 2013, 2014; Midgley & Bond, 2015). Direct impacts of land use change have received less predictive attention (Asner et al., 2010; Heubes et al., 2011; Davies-Barnard et al., 2015) despite the expectation that rapid population growth in tropical Africa (United Nations, 2013), agricultural expansion and intensification (Tilman et al., 2001), and bioenergy production (Alexandratos & Bruinsma, 2012; Laurance et al., 2014) constitute a direct threat both to the biodiversity and the sustainability of services provisioning of savanna and forest ecosystems in the tropics (Sala et al., 2000; Geist & Lambin, 2002; Gardner et al., 2010). Even fewer studies have examined climate change and land use change together (Heubes et al., 2011; Davies-Barnard et al., 2015), and so their relative effects on tropical vegetation remain largely unknown.

Correspondence: Julie C. Aleman, tel. 203 432 5902, fax 203 432 5176, e-mail: julie.aleman@yale.edu

© 2016 John Wiley & Sons Ltd

In sub-Saharan Africa, two biomes dominate: treedominated forests and grass-dominated savannas. Tree cover effectively differentiates the two: forest ecosystems are characterized by a closed tree canopy and a shade-tolerant understory (Ratnam *et al.*, 2011), while savannas are defined by the coexistence of a continuous grass layer with a discontinuous tree layer (Ratnam *et al.*, 2011; Parr *et al.*, 2014). Tree cover therefore differentiates forests from savannas at large scale (Hirota *et al.*, 2011; Staver *et al.*, 2011b).

Savanna and forest distribution and structure result from interactions between climate, soils, and disturbance regimes, especially fire and herbivory (Sankaran *et al.*, 2005). Rainfall is a primary although not singular driver of biome distribution globally (Hirota *et al.*, 2011; Staver *et al.*, 2011b); rainfall constrains maximum tree cover (Sankaran *et al.*, 2005), but tree cover varies substantially below this maximum due to edaphic and top-down processes, including fire and herbivory (Bucini & Hanan, 2007; Bond, 2008; Sankaran *et al.*, 2008). Fire and herbivory themselves represent complex ecological processes: feedbacks with vegetation can strongly influence both fire regimes and herbivore population dynamics (Bond, 2008); abundant grassy fuels make fires vastly more likely in savannas than in forests (Hoffmann *et al.*, 2003), while the accessibility of forage in savannas directly elevates herbivore densities (Riginos & Grace, 2008).

Climate and land-use change will potentially affect each of these processes. In the tropics, climate change is expected to lead to increasing temperature and, more importantly, to changing precipitation patterns (IPCC, 2014). Although fire and herbivory pre-date humans in savannas (Bird & Cali, 1998; Keeley & Rundel, 2005), human activities have substantial impacts on both fire and herbivory regimes globally (Van Langevelde *et al.*, 2003). Land-use, especially agriculture, can also have significant direct effects on ecosystem structure and on tree cover (Laurance *et al.*, 2014). Changes in rainfall, fire, and herbivory regimes, and land-use will thus likely have major impacts on tree cover, of interest in and of themselves from an ecosystems perspective.

However, tree cover change is also tightly linked to habitat loss in both savanna and forest systems, especially when it results from land-use change. A degraded forest is not a savanna, nor is an encroached savanna a forest (Veldman & Putz, 2011; Mitchard & Flintrop, 2013; Veldman et al., 2015a). Indeed, tree cover and biodiversity usually decrease in tropical forests that are logged or subject to intensive agriculture (Barlow et al., 2007; Laurance et al., 2007; Gardner et al., 2009; Gibson et al., 2011). Moreover, degraded forests, where the canopy has been opened resulting in grass invasion, are very different in terms of biodiversity and ecosystem services from old-growth savannas (Veldman & Putz, 2011; Veldman et al., 2015a). Savanna structure is also related to habitat and diversity, both faunal and floral (Du Toit, 1996; Du Toit & Cumming, 1999; Price & Morgan, 2008; Zaloumis & Bond, 2011), and agricultural conversion to intensive cropping and ranching is responsible for tree cover modification and associated biodiversity. However, protected areas have the potential to mitigate against the effects of climate and land-use change on species distributions (Loarie et al., 2009; Ordonez et al., 2014), such that the net effect of tree cover loss on natural and managed systems depends on their conservation context.

Here, we evaluate the potential influences of climate and land-use change on tree cover in tropical Africa. We begin by examining climate, disturbance, and land-use effects on current tree cover, and this relationship to predict the future of tree cover under a variety of different climate and land-use change scenarios. By comparing these expected changes against baseline biome distributions (White, 1983) and protected area distributions from the World Database on Protected Areas, we then evaluate which areas of tropical Africa are potentially most vulnerable to global change.

Material and methods

We use tree cover as a proxy for vegetation structure and land cover (Mayaux *et al.*, 2004) and build a statistical model linking tree cover data in sub-Saharan Africa with climatic, edaphic, and anthropogenic data, from global and remote sensing databases. We use climate and land-use projections from the Representative Concentration Pathways scenarios (Van Vuuren *et al.*, 2011) to simulate tree cover changes in the year 2070. We compare tree cover changes against historical biome distributions (White, 1983) and protected area extent, by WWF Ecoregion [defined as areas sharing homogeneous biophysical features (Olson *et al.*, 2001)], to identify areas of high biodiversity and ecosystem services concern.

Data

Percent tree cover data for the year 2000 were derived from the 250 m MOD44B Collection 5 product from the MODerateresolution Imaging Spectroradiometer (MODIS) sensor (Dimiceli et al., 2011). This product gives percent canopy cover and was calibrated only against trees >5 m tall (Hansen et al., 2003), thus potentially underestimating shrub cover. Rainfall and seasonality are expected to have a greater influence on vegetation structure than temperature in the tropics (Sankaran *et al.*, 2005). Mean annual rainfall (mm yr^{-1}) and seasonality were derived from the 30-arcsec WorldClim Version 1.4 dataset (data averaged from 1950 to 2000, Table S1, Hijmans et al., 2005). The seasonality index used here represents a measure of the variation in monthly rainfall totals over the course of the year and is computed as the ratio of the standard deviation of the monthly total rainfall to the mean monthly total rainfall, expressed as a percentage (Hijmans et al., 2005). The soil map of Africa at 1 km resolution was obtained from the ISRIC World Soil Information team in collaboration with the African Soil Information System project (www.isric.org) and was used to extract soil properties estimates for six soil characteristics that can be potential determinants of savanna structure (Sankaran et al., 2005, 2008; Bucini & Hanan, 2007): percent sand, clay and silt, organic carbon content (g kg⁻¹), pH (~ phosphorus availability) and the cation exchange capacity (~ fertility, in cmol kg^{-1}) average for the top 100 cm of soil (Table S1). We used the monthly L3JRC burnt area product to derive an estimate of fire frequency (Giglio et al., 2010). For this analysis, monthly data layers from 2000 to 2007 were combined to calculate the total number of times individual pixels burned over the time period (Table S1). We used population density data for the year 2000 (Bengtsson et al., 2006). Finally, land-use data (here, the proportion of cropland and pasture per pixel; Table S1), were extracted and computed for the year 2000 from the Harmonized Global Land Use database version 1 (Chini et al., 2014), available for the entire globe at 0.5-degree spatial resolution.

Our analysis was restricted to low elevation and tropical and subtropical climates. Pixels with elevation >1500 m or less than 0 m were excluded using the elevation layer from the Food and Agriculture Organization Harmonized World Soils Database. We also excluded sites with winter rainfall, which are characterized by vegetation other than savanna and forest (Staver *et al.*, 2011b). Finally, we compiled the World Wildlife Fund (WWF) Terrestrial Ecoregions (Olson *et al.*, 2001) and protected areas from the World Database on protected Areas (WDPA) Annual Release 2014 (web download version – February 2015; http://www.protectedplanet.net) for sub-Saharan Africa.

All data were re-projected in WGS-84 at a resolution of 0.08333° (~10 km) using a nearest neighbor procedure. We extracted percent tree cover, fire frequency, climate, soil, and land-use data on a regular grid of 215 160 points for statistical analysis. We computed Pearson correlation coefficients among all predictor variables, and chose only one variable for inclusion in analysis when variables covaried by r > 0.7. We calibrated the model with 70% of the dataset and used the remaining 30% for validation.

Modeling framework

Random forest models use a classification and regression tree approach that recursively partitions predictor variables. The algorithm creates multiple bootstrapped regression trees without pruning and averages the outputs; each tree is grown using a randomized subset of predictors (Breiman, 2001). These models are very effective in reducing variance and error in high dimensional data sets by taking an ensemble of unpruned trees. Moreover, growing large numbers of trees reduces overfitting, and random predictor selection keeps bias low, providing models appropriate for use in prediction (Prasad et al., 2006). Several metrics are available to help interpreting these models. Variable importance can be evaluated based on how much predictions deteriorated when values for the predictors were randomly permuted (Breiman, 2001); we used this approach to compare relative importance among predictors variables. All data were extracted and analyzed in R (R Core Team, 2015), using the "raster" and "randomForest" packages.

Future projections

The emissions scenarios that provide inputs to climate models are produced by Integrated Assessment Models (IAMs), driven by assumptions about socioeconomic forces (e.g. demography, economy, land-use..., Moss *et al.*, 2010). Over time, these scenarios have provided more complex and comprehensive information about air pollutant emissions and land-use (Moss *et al.*, 2010). The latest set of scenarios is contained in four Representative Concentration Pathways (RCPs) which correspond to different radiative forcing trajectories. Each scenario is 'representative' as it provides only one possible trajectory for greenhouse gas and radiative forcing reached by the year 2100 (IPCC, 2014). The term 'pathway' emphasizes that the scenarios represent trajectories over time. RCPs are thus sets of scenarios with independent and clear narratives of emission, socioeconomic and policy trajectories produced by four individual IAMs (Moss *et al.*, 2010).

The RCP 8.5, developed by the MESSAGE modeling team (Riahi et al., 2007), is characterized by increasing greenhouse gas emissions with radiative forcing exceeds 8.5 W m⁻² by 2100 and continues to rise afterward. Within this scenario, a global population increase drives a strong increase in croplands and pasture lands, especially in developing countries (Hurtt et al., 2011). The RCP 6.0 and 4.5, developed, respectively, by the AIM and the GCAM modeling teams (Van Vuuren et al., 2011), correspond to stabilization scenarios wherein radiative forcing stabilizes at ~6 W m⁻² and 4.5 W m⁻² after 2100 due to mitigation actions. In RCP 6.0, croplands are expected to increase due to increasing food demand, but pasture areas somewhat decrease due to a shift from extensive to more intensive husbandry. The RCP 4.5 predicts a radical change in global land-use because carbon from vegetation will be valued as part of global climate policy (Van Vuuren et al., 2011); cropland and pasture thus decrease as a combined result of reforestation programs, yield improvement and dietary changes. Finally, the RCP 2.6, developed by the IMAGE modeling team (Van Vuuren et al., 2011), corresponds to a pathway where radiative forcing peaks at 2.6 W m⁻² before 2100 and then declines. A crucial feature of the RCP 2.6 is the use of bio-energy and carbon capture and storage technologies, which results in negative emissions (Van Vuuren et al., 2011); however, this scenario achieves decreases in radiative forcing via a large increase in croplands dedicated to biofuel production, and pasture also increases as animal production does (Hurtt et al., 2011). In summary, each RCP achieves its radiative forcing trajectories by simulating diverse land-use, socioeconomic and policy scenarios, such that the intensity of land-use change does not monotonically increase with RCP radiative forcing (Van Vuuren et al., 2011).

The Global Circulation Models (GCM) within the CMIP5 framework (Taylor *et al.*, 2012) used the latest release of RCPs as inputs. We thus used an ensemble of all GCM outputs, available downscaled and calibrated against Worldclim 1.4 as baseline climate (Hijmans *et al.*, 2005), for the four RCPs as future precipitation and seasonality projections in 2070 (average 2061–2080). Future land-use projections were derived from the IAM outputs and harmonized against historical and future data (Hurtt *et al.*, 2011). We downloaded cropland and pasture data from Chini *et al.* (2014) and averaged the variables from 2061 to 2080 for the four RCPs for consistency with the climate data. Human population density data were derived from IPPC SRES projections A1B for the year 2100 (SI).

To discriminate between climate vs. land-use change, we calculated future tree cover with the Random Forest model we calibrated and using projected (i) climate change, (ii) land-use change, and (iii) both. As an approximation, we only changed climate, land-use and population variables, keeping fire and soil constant. Indeed, standard future fire projections are not yet available, and changing soil variables would require building a more complex model taking into account feedbacks between soil and vegetation, which is not possible with a statistical model. Moreover, while our future projections do not explicitly consider the alternative stable state dynamic thought to characterize savanna and forest distributions (Hirota *et al.*, 2011; Staver *et al.*, 2011b), preserving present fire patterns across Africa, except where land-use and climate explicitly change tree cover, implies that initial conditions largely determine the extent to which fire feedbacks contribute to determining the distribution of tree cover in tropical biomes (Staver *et al.*, 2011b).

We present results for RCP 2.6 and RCP 4.5 in the main body, because they highlight a strong contrast; RCP 2.6 assumes decreasing CO_2 emissions ultimately, but drastic cropland extent increase in response to biofuel demand, while RCP 4.5 assumes increasing emissions but decreasing cropland and pasture extent through agricultural intensification (Van Vuuren *et al.*, 2011). We include results from the two remaining scenarios (RCP 6.0 and 8.5) in the Supplementary Information. All the model outputs are available to download as raster files at http://www.juliealeman.com/GCB-15-1799/.

Risk assessment

Tree cover is higher in forest than in savanna, therefore the tree cover threshold necessary for vegetation structure to have negative effects on biodiversity and other ecosystem functions, and thus on habitat in general, is thus lower in savanna than in forest. For our analyses, we thus assumed that forest would experience habitat loss when tree cover change decreased by 20% or more, and that savanna, and grassy biomes in general, would experience habitat loss when tree cover changed (either increased or decreased) by more than 10%. Indeed, both decreases in woody cover (Veldman & Putz, 2011) and woody encroachment (Bassett *et al.*, 2000; Mitchard *et al.*, 2009; Wigley *et al.*, 2009; Mitchard & Flintrop, 2013) can have negative impacts on habitat in grassy biomes.

We assessed global change vulnerability by comparing absolute tree cover changes with the abundance of protected areas. The rationale behind this idea is that large tree cover changes, and thus habitat loss, can potentially be buffered by a large network of protected areas (Loarie *et al.*, 2009; Ordonez *et al.*, 2014) in some ecoregions. Protected areas have the potential to mitigate against the effects of climate and land-use change on species distributions (Ordonez *et al.*, 2014), such that the net effect of tree cover loss on natural and managed systems depends on their conservation context.

To maintain a focus on biodiversity responses, we analyzed patterns at the scale of the WWF Terrestrial Ecoregion (Fig. S3), defined as areas with similar floral and faunal communities (Olson *et al.*, 2001). For each future scenario, we plotted the percentage of protected area against the absolute change in tree cover for each ecoregion. For each ecoregion, we computed and plotted the median of the percentage of protected area and the median of absolute tree cover change; ecoregion risk was defined by the quarters of the protected area vs. tree cover change graph, where quarters were demarcated by the median of each variable.

Thus, if an ecoregion had a percentage of protected area below the median and an absolute tree cover change greater than the median, this ecoregion was estimated as 'at risk' (red ecoregion quarter, Figs 3 and 4). Conversely, if an ecoregion had a percentage of protected area above the median and an absolute tree cover change greater than the median, this ecoregion was estimated as at 'mild risk' (pink ecoregion quarter). Thirdly, if an ecoregion had a percentage of protected area above the median but an absolute tree cover change superior to the median, this ecoregion was estimated as at 'no risk' (blue ecoregion quarter). Finally, if an ecoregion had a percentage of protected area below the median and an absolute tree cover change inferior to the median, this ecoregion was estimated as at 'no risk but to be monitored' (light blue ecoregion quarter).

Results

Tree cover modeling and determinants

The model predicts tree cover with high accuracy (Fig. 1b, $R^2 = 0.97$, $N = 20\ 000$, P < 0.01, *t*-test), and also captured the spatial pattern of forest and savanna distributions well. Indeed, only 3.7 and 0.7% of forest



Fig. 1 Model predictor importance (a) and model evaluation (b). MAR = mean annual rainfall; CEC = Cation exchange capacity. Relative importance of model predictors was computed by randomly permuting variables and evaluating model deterioration. Model evaluation was performed on a subset of tree cover pixels in Africa ($N = 20\ 000$, $R^2 = 0.97$, RMSE = 4.47).



Fig. 2 Tree cover and biome distribution in tropical Africa. Tree cover for the year 2000 from MODIS data (a), predicted tree cover for the year 2000 using the Random Forest model (b), the difference between real and modeled tree cover (c), and biome classification/misclassification (d). Forest and savanna biomes were defined using a tree cover threshold of 55 (Staver *et al.*, 2011b).

and savanna pixels, respectively, were misclassified (Fig. 2a–d), and these were generally restricted to locations near savanna-forest boundaries. While our model does not consider the alternative stable state dynamic, thought to characterize savanna and forest distributions (Hirota *et al.*, 2011; Staver *et al.*, 2011b), it clearly captures, at least correlatively, the bimodal nature of tree cover in sub-Saharan Africa (Fig. S1).

Climatic variables were strong determinants of tree cover (Fig. 1a), highlighting the importance of annual rainfall and seasonality in determining forest and savanna distributions. Tree cover showed a strong positive dependence on annual rainfall between 0 and 2000 mm and a negative one with seasonality (Fig. S2). Anthropogenic variables also mattered, especially pasture coverage, through pastoralism, which has a very strong negative correlation with tree cover (Figs 1a and S2). Cropland intensity in pixels also reduces tree cover and the extent both of forest and savanna (Fig. S2). As expected, fire has major impacts on tree cover, with decreasing tree cover by increasing fire frequency (Fig. S2). Finally, and not surprisingly at this scale, edaphic variables were significant, but less predictive of tree cover; soil organic carbon concentration was the most predictive soil variable (Fig. S2), potentially reflecting vegetation impacts on soil carbon, and not vice versa.

Future tree cover and habitat loss projections

Forests experienced significant tree cover changes in all simulations throughout tropical Africa (Figs 3, 4, S4 and S5). Under RCP 2.6, tree cover decreased drastically in the forest areas inducing up to ~27% of habitat loss (Fig. 3). Indeed, the Congo forest contracted (especially in Cameroon, Central African Republic, Guinea, Gabon and Uganda) and fragmented (especially in the Democratic Republic of Congo, hereafter DRC, and northern Republic of Congo) (Fig. 3; Table S2). Tree cover decreased provoking the loss of habitat in up to 76.6% of the Guinean forest block (Liberia and Sierra Leone), and to 96.7% of dry forests throughout Africa (in Madagascar, Ethiopia, Angola and Mozambique; RCP 2.6, Table S2). Tree cover in forests also mostly decreased, although somewhat less and with localized increases, under RCP 4.5 (Fig. 4 and Table S2). Interestingly, these widespread changes in tree cover occurred primarily because of changes in pasture and cropland extent (Figs 3b-c, 4b-c, S8 and S9); agricultural extensification under RCP 2.6 drove decreases in tree cover, while agricultural intensification (and concomitant contraction) resulted in some tree cover increases in forests under RCP 4.5. Despite a general increase in seasonality with RCPs (Fig. S7), climate did not drive tree cover changes at the continental scale,

6 J. C. ALEMAN et al.

(a) Global change



Fig. 3 Tree cover change projections from 2000 to 2070 – RCP 2.6 (left maps), biome distribution shifts based on tree cover changes (middle), and WWF Ecoregion conservation risk based on tree cover change and protected area coverage (right). Projections are based on climate and anthropogenic change together (a), anthropogenic change only (b), and climate change only (c).

even under the most extreme climate scenario (RCP 8.5; Fig. S5).

Meanwhile, tree cover was also projected to decrease in savanna throughout tropical Africa (Figs 3, 4, S4 and S5), but trajectories varied geographically and by scenario. Tree cover in savannas generally decreased under RCP 2.6, especially in savanna areas bordering forests (Fig. 3). Under RCP 4.5, savanna tree cover decreased in southern and some parts of Central Africa (DRC, Cameroon, Angola, Zambia, Mozambique), but increased elsewhere (West and littoral Central Africa and Madagascar; Table S2 and Fig. 4). Comparing projected increases in tree cover with a reclassified map of vegetation types in Africa adapted from White's (1983) phytogeographic map (Fig. S3c), we identified that 36.8% (and 40.0% for RCP 6.0) of tree cover increases in mesic savannas actually represent regrowth of forests deforested between 1983 and 2000. However, the remainder (43.2% and 60.0% for RCP 4.5 and 6.0, respectively) represents woody encroachment of mesic savanna. Again, tree cover changed mostly as a result of changing cropland and pasture extent (Figs S8 and

(a) Global change



Fig. 4 Tree cover change projections from 2000 to 2070 – RCP 4.5 (left maps), biome distribution shifts based on tree cover changes (middle), and WWF Ecoregion conservation risk based on tree cover change and protected area coverage (right). Projections are based on climate and anthropogenic change together (a), anthropogenic change only (b), and climate change only (c).

S9), not as a consequence of climate change, despite a predicted increase in seasonality in some areas (Fig. S7).

The forest areas most vulnerable to habitat loss in our predictions were also associated with a large network of protected areas (Figs 3 and 4). However, in savannas, where tree cover also changed substantially, if less consistently, the extent of protected area coverage varied more substantially (Fig. S3b). Under RCP 2.6, tree cover in Zambia, Mozambique and Tanzania (Fig. 3) decreased strongly, but these areas were generally well-protected. Tree cover in other savannas (*e.g.*

bordering forests and in Angola; Fig. 3) also decreased under RCP 2.6, but protected areas were limited in extent.

Discussion

Vegetation structure and tree cover determinants

The model we developed here predicts tree cover very well, especially compared to most of other attempts in predicting tree cover at the continental scale (Sankaran *et al.*, 2005; Bucini & Hanan, 2007; Staver *et al.*, 2011a),

probably because we included land-use variables as predictor. These are known to play an important role in determining vegetation structure currently (Veldman et al., 2015a), especially in savanna ecosystems where land-use has potentially shaped land-cover for millennia (Kay & Kaplan, 2015; Veldman et al., 2015a). Although pastoralism has minimal impact on vegetation cover [e.g., in Central African Republic (Ankogui-Mpoko, 2003)] and can even lead to bush encroachment (Bassett et al., 2000), recent land-use intensification has changed vegetation structure associated with significant biodiversity decrease (Foley et al., 2005). Shifting agriculture in the tropics consists of deforested small patches of forest or savanna for creating cropland, that are then transformed into fallows, obviously decreasing tree cover (Nacoulma et al., 2011). Once again, intensive agriculture increases the rate of tree removal (Laurance et al., 2014).

The importance of climatic variables in determining tree cover was not surprising, since climate constitutes the primary driver for tree cover in Africa (Sankaran *et al.*, 2005, 2008; Bucini & Hanan, 2007). Indeed, annual rainfall determines the climatic envelop where only savanna, savanna and forest, and finally only forest are possible (Staver *et al.*, 2011a). Seasonality and thus water availability is known to be a strong determinant for tree cover in Africa (Bucini & Hanan, 2007; Sankaran *et al.*, 2008; Good & Caylor, 2011), both via direct effects on trees and grasses growth rates (Sankaran *et al.*, 2004) and on the probability of fire spread.

Still, some pixels were misclassified, especially in areas of transitions between the two biomes. This is a well-known issue (Zeng *et al.*, 2014) and probably occurs because forest and savanna are not deterministically related to climate (Staver *et al.*, 2011a). We did not consider the alternative stable state dynamic of savanna and forest distributions (Hirota *et al.*, 2011; Staver *et al.*, 2011b), in part because the model is correlative, linking variables statistically and not dynamically, but also because land-use-driven transitions are not part of the positive feedback that maintains savanna and forest as stable states (Staver *et al.*, 2011a).

Projected changes and the vulnerability of tropical ecosystems

Despite a projected general increase in seasonality, we showed here that climate was not the primary driver of tree cover changes at the continental scale, regardless of scenario and even under the most extreme climate change. This contradicts previous studies that have suggested that changes in vegetation structure would be primarily precipitation-driven (Scheiter & Higgins, 2009; Good & Caylor, 2011; Zeng *et al.*, 2014).

Instead, we showed that land-use change was the main driver of tree cover changes. This should not be wholly surprising, even in savannas, which are often perceived as more anthropogenic than forests (Parr *et al.*, 2014); recent studies have shown that bigger land cover changes originated from land-use than from climate change on shorter time scales (Arneth, 2015; Davies-Barnard *et al.*, 2015), and biodiversity projections have suggested that land-use change can lead to drastic erosion of vegetation structure and biodiversity (Sala *et al.*, 2000; Asner *et al.*, 2010).

Tree cover decreases in forest (>20%) resulting from land-use intensification will be extensive under all scenarios. Under 'mild' emissions scenarios (RCP 2.6), agricultural intensification will result from biofuel production while under more severe scenarios (RCP 8.5) it will result from food crops production mostly for feeding increasing human populations. Impacts on plant and animal diversity as well as ecosystem services will then likely be extensive (Laurance et al., 2014). Indeed, "old-growth" forests are irreplaceable (Barlow et al., 2007; Gibson et al., 2011) in terms of biodiversity, and studies have shown that forest biodiversity declines along a gradient of land-use (Schulze et al. 2004; Harvey et al. 2006; Basset et al. 2008; Philpott et al. 2008), reflecting the decline in floristic and structural diversity. Forest fragmentation is causing high rates of extinction (Turner, 1996). Moreover, deforestation can cause loss of topsoil and alteration of the water flow, resulting in either flood or drought (Bradshaw et al., 2008). It affects vertebrates' habitat and so reduces their diversity (Jetz et al., 2007; Nichols et al., 2007; Slade et al., 2011). Finally, secondary forests and plantation are much less diverse than primary forests (Barlow et al., 2007; Laurance, 2007).

Likewise, the replacement of "old-growth" grassy biome by grass-dominated "secondary" vegetation is often associated with a plant diversity collapse (Veldman & Putz, 2011). Land-use intensification will have thus equally severe impacts in grass-dominated biomes. The Brazilian cerrado, for instance, is highly threatened as a result of large-scale conversion mainly to soybean and cattle ranching, resulting in water pollution, fire regime modifications, and biodiversity loss (Klink & Machado, 2005). The degradation of grassy biomes by agricultural conversion results in the displacement of native species, the alteration of soil chemistry, fire regime, and hydrology (Veldman et al., 2015a). Therefore, as in forests, "secondary" grassy ecosystems are less diverse and resilient than the "oldgrowth" ones (Zaloumis & Bond, 2011; Veldman et al., 2015a).

In our simulations, humid savannas were highly threatened for RCPs 2.6 and 8.5, in large part because

they have been identified as suitable for biofuel production and agriculture intensification (Deininger & Byerlee, 2011; Alexandratos & Bruinsma, 2012). These studies mistakenly (Parr et al., 2014; Veldman et al., 2015a) assume savannas to be less valuable than forests in terms of biodiversity, carbon storage, and water quality, and consequently meriting less protection (Searchinger et al., 2015; Veldman et al., 2015b). Converting these savannas to agricultural lands would have catastrophic consequences for numerous endemic species and the high regional bird, mammal, and flora richness they sustain (Searchinger et al., 2015). Grassdominated ecosystems contain numerous hotspots of endemism and the conversion of those to agricultural lands will inevitably lead to irreplaceable species loss (Olson et al., 2001). Once again consider the example of the Brazilian cerrado, which is still considered of less conservation value than tropical forest; soybean production and ranching have reduced native savanna extent by more than 50% (Klink & Machado, 2005). A switch in land-use practices, in African humid savannas, from shifting cultivation to cash crops and biofuel production will increase the pressure of agricultural systems on the vegetation, with potential consequences similar to what is currently happening in the *cerrado*. Moreover, switching from extensive agriculture to a more intensive one, oriented in cash-crops and biofuel production (RCP 2.6), raises food security questions (Escobar *et al.*, 2009)

On the other hand, simulations for RCPs 4.5 and 6.0 project an increase in tree cover in savanna areas; some of this corresponds to forest regrowth after agricultural abandonment, but the majority represents forest encroachment of savanna, with major potential negative consequences. Woody encroachment reduces plant species richness and locally modifies the germinable seed bank (Meik *et al.*, 2002; Price & Morgan, 2008; Sirami *et al.*, 2009). It is also responsible for changes in the composition and spatial distribution of vegetation structures (Skarpe, 1986; Jeltsch *et al.*, 1997), which are accompanied by changes in availability and accessibility of resources (e.g. foraging sites or predation cover), thus affecting rodent communities (Blaum *et al.*, 2007a) and mammalian carnivores (Blaum *et al.*, 2007b).

Protected areas and buffering habitat loss

In the face of massive global changes, protected areas are crucial to sustaining biodiversity and natural ecosystem processes. In forests, where our model predicted widespread reductions in tree cover, a large network of protected areas already potentially confers some degree of local resilience to change. However, concern is widespread that protected areas are not

effectively enforced, due primarily to conflict and governance issues (Chape et al., 2005), potentially compromising their role in maintaining forest resilience. Indeed, tropical protected areas face growing threats as population grows (Laurance et al., 2014), especially due to limited funding for management (Bruner et al., 2004) and illegal logging, grazing, and agricultural lands are common in poorly enforced protected areas (Laurance et al., 2012), especially in West and Central Africa (Tranquilli et al., 2014) compared to East and South Africa (Pfeifer et al., 2012). As a result, agricultural expansion near protected areas tends to erode biodiversity due to edge effects (Wittemyer et al., 2008; Laurance et al., 2014). However, if protected area integrity can be achieved, as for example by involving local population (Vodouhê et al., 2010), African forests may remain relatively resilient in the face of land-use and rainfall change.

By contrast, savannas where tree cover change is likely are relatively unprotected. For example, we projected savannas from Zambia and southern DRC, which currently sustain some of the world's highest terrestrial mammal species relative richness (Olson et al., 2001), to experience drastic tree cover reductions (RCP 2.6). However, while Zambian savannas are extensively protected, savannas in the southern DRC are not, such that the consequences of decreasing tree cover potentially diverge. One possibility solution is an increase in formal conservation activities (parks and otherwise) within savanna ecosystems, especially in West and Central Africa, which may mitigate or even prevent tree cover effects of land-use change in the future (Parr et al., 2014; Veldman et al., 2015b). Grassy biomes may also lend themselves to alternative conservation practices. Protected areas in grassy biomes are often associated with reduced plant diversity compared to communally managed areas (Dahlberg, 2000; Shackleton, 2000; Hahn-Hadjali et al., 2006; Paré et al., 2010; Nacoulma et al., 2011), perhaps because grassy biomes are intrinsically associated with disturbances (Veldman et al., 2015a), such that traditional land management does not lead to degradation of savanna habitats (Augusseau et al., 2006; Nacoulma et al., 2011). Indeed, re-evaluation of systematic landuse change planning for mitigations scenarios may also be warranted, to explicitly include the conservation and agricultural value of savanna ecosystems (Veldman et al., 2015a). Some authors have even argued that effective biodiversity conservation will rely on associating traditional, communally managed areas with reserves (Abel & Blaikie, 1989; Nacoulma et al., 2011; Yayneshet & Treydte, 2015). The status for grassy biomes conservation in protected areas is currently too linked to management of forest ecosystems.

There is thus an urgent need to develop a framework for grass-dominated biomes (Veldman *et al.*, 2015a,b) and to better estimate the environmental values of such biomes.

Historically, the vast majority of conservation work has focused on forests (Laurance *et al.*, 2014; Tranquilli *et al.*, 2014), but savannas are at equal if not greater risk of change under plausible land-use change scenarios. Wet savannas and transition zones between forest and savannas in West and Central Africa may be particularly susceptible, because of the few protected areas located there. Sustainable management plans, including a strong monitoring component, serving both biodiversity and development needs, should be a high priority.

Limitations of the approach used

Here, we explicitly considered neither direct CO₂ nor temperature effects on vegetation, although under RCP 8.5, $[CO_2]$ more than doubles and temperatures could rise locally by as much as 4 °C in tropical Africa. Rising [CO₂] may promote woody encroachment in increasingly arid environments by increasing growth rates and water use efficiency (Bond & Midgley, 2012), such that increasing CO₂ could result in shifts from savanna toward a woodier plant community, such as thicket or even closed forest (Higgins & Scheiter, 2012). We were unlikely to detect this possibility in any case, since the MODIS VCF does not record trees with less than 5 m height and therefore only poorly captures the conservation and functional costs of woody encroachment, which are ongoing in savannas worldwide. These studies still do not take into account land-use (Midgley & Bond, 2015), however, which may be at least locally sufficient to counter CO₂-related changes (Sala et al., 2000; Midgley et al., 2010; Wessels et al., 2013; Tredennick & Hanan, 2015).

The extant range of temperature variability projected for the future exceeds the current one, such that our statistical model is not built to take it into account. Although temperature changes are expected to be less severe than in boreal and temperate ecosystems (IPCC, 2014), rising temperatures could have major negative effects on tropical vegetation, especially forest trees (Allen et al., 2010), and are likely to exacerbate the consequences of land-use change on tree cover. Additionally, another source of discrepancy in our simulations may originate from the high uncertainty in future climate projections that are related to GCM biases (Knutti & Sedláček, 2013) and which are especially noticeable in Africa (Heubes et al., 2011). However, our conclusions regarding land-use changes being the main driver of vegetation structure modifications in the future remain supported because the scenarios are coherent regarding cropland and pasture projections (Hurtt *et al.*, 2011).

Finally, fire has a key role globally in the distribution of savanna and forest by maintaining a feedback on vegetation structure (Staver *et al.*, 2011a) and locally to determine savanna tree cover (Nepstad *et al.*, 1999; Veldman & Putz, 2011; Brando *et al.*, 2014). Future changes in fire regimes may therefore impact the structure of both forest and savanna (Nepstad *et al.*, 1999; Veldman & Putz, 2011; Brando *et al.*, 2014). Unfortunately, global fire models are still in their nascency, and no standardized projections for future fire regime are currently available. Incorporating fire model projections into biosphere projections for tropical and sub-tropical Africa will be critical, once they are more available.

To conclude, we evaluated the potential contributions of climate and land-use changes to changing tree cover in tropical Africa, and the implications for potential habitat loss. Land-use changes were the main driver of tree cover change. Moreover, we show that savannas are at an equal if not greater risk of change under global change scenarios. Finally, our results also highlight the importance of how global and local policies decide to mitigate climate change. Mitigation scenarios by definition overwhelmingly prioritize a reduction in carbon emissions, but at what cost? Biofuel production and agricultural intensification potentially result in direct negative consequences for forest and especially savanna extent in tropical Africa.

Acknowledgements

We thank two anonymous reviewers for their valuable comments and suggestions that greatly improved our manuscript. Funding for this work was provided by the Forest and Elizabeth Brown Fund for postdoctoral fellowships in the plant sciences at Yale University. Data were obtained from the ISRIC - World Soil Information team, Worldclim and WWF team, and we gratefully acknowledge the work of the data contributors and the community.

References

- Abel N, Blaikie P (1989) Land degradation, stocking rates and conservation policies in the communal rangelands of Botswana and Zimbabwe. Land Degradation & Development, 1, 101–123.
- Alexandratos N, Bruinsma J (2012) World agriculture towards 2030/2050: the 2012 revision. ESA Work. Pap, 3.
- Allen CD, Macalady AK, Chenchouni H et al. (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259, 660–684.
- Ankogui-Mpoko GF (2003) Gestion des espaces agropastoraux en zones de savanes centrafricaines: De l'arbitrage de l'Etat à une cogestion. In: Savanes Africaines: Des Espaces en Mutation, des Acteurs Face à de Nouveaux Défis (eds Jamin JYSBL, Floret C). Prasac, N'Djamena, Tchad - Cirad, Garoua, Cameroun, Montpellier, France.
- Arneth A (2015) Climate science: Uncertain future for vegetation cover. Nature, 524, 44-45.
- Asner GP, Loarie SR, Heyder U (2010) Combined effects of climate and land-use change on the future of humid tropical forests. *Conservation Letters*, **3**, 395–403.

- Augusseau X, Nikiéma P, Torquebiau E (2006) Tree biodiversity, land dynamics and farmers' strategies on the agricultural frontier of southwestern Burkina Faso. *Biodi*versity & Conservation, 15, 613–630.
- Barlow J, Gardner TA, Araujo IS et al. (2007) Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. Proceedings of the National Academy of Sciences of the United States of America, 104, 18555–18560.
- Basset Y, Missa O, Alonso A et al. (2008) Changes in arthropod assemblages along a wide gradient of disturbance in Gabon. Conservation Biology, 22, 1552–1563.
- Bassett TJ, Boutrais J, Cline-Cole R, Madge C (2000) Cattle and trees in the West African savanna. In: *Contesting Forestry in West Africa* (eds Cline-Cole R, Madge C), pp. 242–263. Ashgate Publishing Ltd, Aldershot.
- Bengtsson M, Shen Y, Oki T (2006) A SRES-based gridded global population dataset for 1990–2100. Population and Environment, 28, 113–131.
- Bird M, Cali J (1998) A million-year record of fire in sub-Saharan Africa. Nature, 394, 767–769.
- Blaum N, Rossmanith E, Jeltsch F (2007a) Land use affects rodent communities in Kalahari savannah rangelands. African Journal of Ecology, 45, 189–195.
- Blaum N, Rossmanith E, Popp A, Jeltsch F (2007b) Shrub encroachment affects mammalian carnivore abundance and species richness in semiarid rangelands. Acta Oecologica, 31, 86–92.
- Bond W (2008) What limits trees in C4 grasslands and savannas? Annual Review of Ecology, Evolution, and Systematics, 39, 641–659.
- Bond WJ, Midgley GF (2012) Carbon dioxide and the uneasy interactions of trees and savannah grasses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 601–612.
- Bradshaw CJ, Sodhi NS, Brook BW (2008) Tropical turmoil: a biodiversity tragedy in progress. Frontiers in Ecology and the Environment, 7, 79–87.
- Brando PM, Balch JK, Nepstad DC et al. (2014) Abrupt increases in Amazonian tree mortality due to drought–fire interactions. Proceedings of the National Academy of Sciences of the United States of America, 111, 6347–6352.
- Breiman L (2001) Random forests. Machine Learning, 45, 5-32.
- Bruner AG, Gullison RE, Balmford A (2004) Financial costs and shortfalls of managing and expanding protected-area systems in developing countries. *BioScience*, 54, 1119–1126.
- Bucini G, Hanan NP (2007) A continental-scale analysis of tree cover in African savannas. Global Ecology and Biogeography, 16, 593–605.
- Chape S, Harrison J, Spalding M, Lysenko I (2005) Measuring the extent and effectiveness of protected areas as an indicator for meeting global biodiversity targets. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360, 443–455.
- Chini LP, Hurtt GC, Frolking S (2014) Harmonized Global Land Use for Years 1500 2100, V1. Data set. Available at [http://daac.ornl.gov] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA. http://dx.doi.org/10.3334/ORNLDAAC/1248 (accessed 15 February 2015).
- Dahlberg AC (2000) Vegetation diversity and change in relation to land use, soil and rainfall—a case study from North-East District, Botswana. *Journal of Arid Environ*ments, 44, 19–40.
- Davies-Barnard T, Valdes P, Singarayer J, Wiltshire A, Jones C (2015) Quantifying the relative importance of land cover change from climate and land-use in the representative concentration pathways. *Global Biogeochemical Cycles*, 29, 842–853.
- Deininger KW, Byerlee D (2011) Rising Global Interest in Farmland: Can it Yield Sustainable and Equitable Benefits? World Bank Publications, Washington, DC.
- Dimiceli CM, Carroll ML, Sohlberg RA, Huang C, Hansen M, Townshend JRG (2011) Annual Global Automated MODIS Vegetation Continuous Fields (MOD44B) at 250 m Spatial Resolution for Data Years Beginning Day 65, 2000 – 2010, Collection 5 Percent Tree Cover. University of Maryland, College Park, MD.
- Du Toit J (1996) Determinants of the composition and distribution of wildlife communities in Southern Africa. *Biological Conservation*, **2**, 217.
- Du Toit J, Cumming D (1999) Functional significance of ungulate diversity in African savannas and the ecological implications of the spread of pastoralism. *Biodiversity & Conservation*, **8**, 1643–1661.
- Escobar JC, Lora ES, Venturini OJ, Yáñez EE, Castillo EF, Almazan O (2009) Biofuels: environment, technology and food security. *Renewable and Sustainable Energy Reviews*, 13, 1275–1287.
- Foley JA, Defries R, Asner GP *et al.* (2005) Global consequences of land use. *Science*, **309**, 570–574.
- Gardner TA, Barlow J, Chazdon R, Ewers RM, Harvey CA, Peres CA, Sodhi NS (2009) Prospects for tropical forest biodiversity in a human-modified world. *Ecology Letters*, **12**, 561–582.
- Gardner TA, Barlow J, Sodhi NS, Peres CA (2010) A multi-region assessment of tropical forest biodiversity in a human-modified world. *Biological Conservation*, 143, 2293–2300.

- Geist HJ, Lambin EF (2002) Proximate Causes and Underlying Driving Forces of Tropical Deforestation Tropical forests are disappearing as the result of many pressures, both local and regional, acting in various combinations in different geographical locations. *BioScience*, 52, 143–150.
- Gibson L, Lee TM, Koh LP et al. (2011) Primary forests are irreplaceable for sustaining tropical biodiversity. Nature, 478, 378–381.
- Giglio L, Randerson JT, Van Der Werf GR, Kasibhatla PS, Collatz GJ, Morton DC, Defries RS (2010) Assessing variability and long-term trends in burned area by merging multiple satellite fire products. *Biogeosciences*, 7, 1171–1186.
- Good SP, Caylor KK (2011) Climatological determinants of woody cover in Africa. Proceedings of the National Academy of Sciences of the United States of America, 108, 4902–4907.
- Hahn-Hadjali K, Schmidt M, Thiombiano A (2006) Phytodiversity dynamics in pastured and protected West African savannas. In: Taxonomy and ecology of African Plants: Their conservation and sustainable use—Proceedings of the 17th AETFAT Congress Addis Ababa.
- Hansen M, Defries R, Townshend J, Carroll M, Dimiceli C, Sohlberg R (2003) Global percent tree cover at a spatial resolution of 500 meters: First results of the MODIS vegetation continuous fields algorithm. *Earth Interactions*, 7, 1–15.
- Harvey CA, Medina A, Sanchez DM *et al.* (2006) Patterns of animal diversity in different forms of tree cover in agricultural landscapes. *Ecological Applications*, 16, 1986– 1999.
- Heubes J, Kühn I, König K, Wittig R, Zizka G, Hahn K (2011) Modelling biome shifts and tree cover change for 2050 in West Africa. *Journal of Biogeography*, 38, 2248– 2258.
- Higgins SI, Scheiter S (2012) Atmospheric CO2 forces abrupt vegetation shifts locally, but not globally. *Nature*, 488, 209–212.
- Hijmans R, Cameron S, Parra J, Jones P, Jarvis A (2005) WorldClim—global climate data. Very High Resolution Interpolated Climate Surfaces for Global Land Areas.
- Hirota M, Holmgren M, Van Nes EH, Scheffer M (2011) Global resilience of tropical forest and savanna to critical transitions. *Science*, 334, 232–235.
- Hoffmann W, Orthen B, Do Nasciment P (2003) Comparative fire ecology of tropical savanna and forest trees. *Functional Ecology*, **17**, 720–726.
- Hurtt G, Chini LP, Frolking S et al. (2011) Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change*, **109**, 117– 161.
- IPCC (2014) Climate Change 2014: Impacts, Adaptation, and Vulnerability. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Jeltsch F, Milton SJ, Dean WRJ, Van Rooyen N (1997) Analysing shrub encroachment in the southern Kalahari: a grid-based modelling approach. *Journal of Applied Ecol*ogy, 34, 1497–1508.
- Jetz W, Wilcove DS, Dobson AP (2007) Projected impacts of climate and land-use change on the global diversity of birds. PLoS Biology, 5, e157.
- Kay AU, Kaplan JO (2015) Human subsistence and land use in sub-Saharan Africa, 1000 BC to AD 1500: A review, quantification, and classification. *Anthropocene*, 9, 14–32.
- Keeley JE, Rundel PW (2005) Fire and the Miocene expansion of C4 grasslands. Ecology Letters, 8, 683–690.
- Klink CA, Machado RB (2005) Conservation of the Brazilian cerrado. Conservation Biology, 19, 707–713.
- Knutti R, Sedláček J (2013) Robustness and uncertainties in the new CMIP5 climate model projections. Nature Climate Change, 3, 369–373.
- Laurance WF (2007) Have we overstated the tropical biodiversity crisis? Trends in Ecology & Evolution, 22, 65–70.
- Laurance WF, Nascimento HE, Laurance SG et al. (2007) Habitat fragmentation, variable edge effects, and the landscape-divergence hypothesis. PLoS ONE, 2, e1017.
- Laurance WF, Useche DC, Rendeiro J et al. (2012) Averting biodiversity collapse in tropical forest protected areas. Nature, 489, 290–294.
- Laurance WF, Sayer J, Cassman KG (2014) Agricultural expansion and its impacts on tropical nature. *Trends in Ecology & Evolution*, 29, 107–116.
- Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of climate change. *Nature*, 462, 1052–1055.
- Mayaux P, Bartholomé E, Fritz S, Belward A (2004) A new land cover map of Africa for the year 2000. *Journal of Biogeography*, **31**, 861–877.
- Meik JM, Jeo RM, Mendelson Iii JR, Jenks KE (2002) Effects of bush encroachment on an assemblage of diurnal lizard species in central Namibia. *Biological Conservation*, 106, 29–36.
- Midgley GF, Bond WJ (2015) Future of African terrestrial biodiversity and ecosystems under anthropogenic climate change. *Nature Climate Change*, **5**, 823–829.

12 J. C. ALEMAN et al.

- Midgley JJ, Lawes MJ, Chamaillé-Jammes S (2010) Savanna woody plant dynamics: the role of fire and herbivory, separately and synergistically. *Australian Journal of Botany*, 58, 1–11.
- Mitchard ET, Flintrop CM (2013) Woody encroachment and forest degradation in sub-Saharan Africa's woodlands and savannas 1982–2006. Philosophical Transactions of the Royal Society B: Biological Sciences, 368, 20120406.
- Mitchard E, Saatchi S, Gerard F, Lewis S, Meir P (2009) Measuring woody encroachment along a forest-savanna boundary in Central Africa. *Earth Interactions*, 13, 1–29.
- Moss RH, Edmonds JA, Hibbard KA et al. (2010) The next generation of scenarios for climate change research and assessment. Nature, 463, 747–756.
- Nacoulma BMI, Schumann K, Traoré S, Bernhardt-Römermann M, Hahn K, Wittig R, Thiombiano A (2011) Impacts of land-use on West African savanna vegetation: a comparison between protected and communal area in Burkina Faso. *Biodiversity* and Conservation, 20, 3341–3362.
- Nepstad DC, Verssimo A, Alencar A et al. (1999) Large-scale impoverishment of Amazonian forests by logging and fire. Nature, 398, 505–508.
- Nichols E, Larsen T, Spector S et al. (2007) Global dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. *Biological Conservation*, **137**, 1–19.
- Olson DM, Dinerstein E, Wikramanayake ED et al. (2001) Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. BioScience, 51, 933–938.
- Ordonez A, Martinuzzi S, Radeloff VC, Williams JW (2014) Combined speeds of climate and land-use change of the conterminous US until 2050. *Nature Climate Change*, 4, 811–816.
- Paré S, Tigabu M, Savadogo P, Odén PC, Ouadba JM (2010) Does designation of protected areas ensure conservation of tree diversity in the Sudanian dry forest of Burkina Faso? African Journal of Ecology, 48, 347–360.
- Parr CL, Lehmann CE, Bond WJ, Hoffmann WA, Andersen AN (2014) Tropical grassy biomes: misunderstood, neglected, and under threat. *Trends in Ecology & Evolution*, 29, 205–213.
- Pfeifer M, Burgess ND, Swetnam RD, Platts PJ, Willcock S, Marchant R (2012) Protected areas: mixed success in conserving East Africa's evergreen forests. *PLoS ONE*, 7, e39337.
- Philpott SM, Arendt WJ, Armbrecht I et al. (2008) Biodiversity loss in Latin American coffee landscapes: review of the evidence on ants, birds, and trees. Conservation Biology, 22, 1093–1105.
- Prasad AM, Iverson LR, Liaw A (2006) Newer classification and regression tree techniques: bagging and random forests for ecological prediction. *Ecosystems*, 9, 181–199.
- Price JN, Morgan JW (2008) Woody plant encroachment reduces species richness of herb-rich woodlands in southern Australia. Austral Ecology, 33, 278–289.
- R Core Team (2015) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ratnam J, Bond WJ, Fensham RJ et al. (2011) When is a 'forest'a savanna, and why does it matter? Global Ecology and Biogeography, 20, 653–660.
- Riahi K, Grübler A, Nakicenovic N (2007) Scenarios of long-term socio-economic and environmental development under climate stabilization. *Technological Forecasting* and Social Change, 74, 887–935.
- Riginos C, Grace JB (2008) Savanna tree density, herbivores, and the herbaceous community: bottom-up vs. top-down effects. *Ecology*, 89, 2228–2238.
- Sala O, Chapin Iii F, Armesto J *et al.* (2000) Global biodiversity scenarios for the year 2100. *Science*, **287**, 1770.
- Sankaran M, Ratnam J, Hanan NP (2004) Tree-grass coexistence in savannas revisited - insights from an examination of assumptions and mechanisms invoked in existing models. *Ecology Letters*, 7, 480–490.
- Sankaran M, Hanan NP, Scholes RJ et al. (2005) Determinants of woody cover in African savannas. Nature, 438, 846–849.
- Sankaran M, Ratnam J, Hanan N (2008) Woody cover in African savannas: the role of resources, fire and herbivory. Global Ecology and Biogeography, 17, 236–245.
- Scheiter S, Higgins SI (2009) Impacts of climate change on the vegetation of Africa: an adaptive dynamic vegetation modelling approach. *Global Change Biology*, 15, 2224–2246.
- Schulze CH, Waltert M, Kessler PJA et al. (2004) Biodiversity indicator groups of tropical land-use systems: comparing plants, birds, and insects. *Ecological Applications*, 14, 1321–1333.
- Searchinger TD, Estes L, Thornton PK et al. (2015) High carbon and biodiversity costs from converting Africa's wet savannahs to cropland. Nature Climate Change, 5, 481–486.

- Shackleton CM (2000) Comparison of plant diversity in protected and communal lands in the Bushbuckridge lowveld savanna, South Africa. *Biological Conservation*, 94, 273–285.
- Sirami C, Seymour C, Midgley G, Barnard P (2009) The impact of shrub encroachment on savanna bird diversity from local to regional scale. *Diversity and Distributions*, **15**, 948–957.
- Skarpe C (1986) Plant community structure in relation to grazing and environmental changes along a north-south transect in the western Kalahari. *Vegetatio*, 68, 3–18.
- Slade EM, Mann DJ, Lewis OT (2011) Biodiversity and ecosystem function of tropical forest dung beetles under contrasting logging regimes. *Biological Conservation*, 144, 166–174.
- Staver AC, Archibald S, Levin S (2011a) Tree cover in sub-Saharan Africa: rainfall and fire constrain forest and savanna as alternative stable states. *Ecology*, 92, 1063–1072.
- Staver AC, Archibald S, Levin SA (2011b) The global extent and determinants of savanna and forest as alternative biome states. *Science*, 334, 230–232.
- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. Bulletin of the American Meteorological Society, 93, 485–498.
- Tilman D, Fargione J, Wolff B et al. (2001) Forecasting agriculturally driven global environmental change. Science, 292, 281–284.
- Tranquilli S, Abedi-Lartey M, Abernethy K et al. (2014) Protected Areas in Tropical Africa: Assessing Threats and Conservation Activities. PLoS ONE, 9, e114154.
- Tredennick AT, Hanan NP (2015) Effects of Tree Harvest on the Stable-State Dynamics of Savanna and Forest. *The American Naturalist*, **185**, E153–E165.
- Turner I (1996) Species loss in fragments of tropical rain forest: a review of the evidence. Journal of Applied Ecology, 33, 200–209.
- United Nations (2013) World Population Prospects: The 2012 Revision. United Nations, Department of Economic and Social Affairs, Population Division, New York, NY.
- Van Langevelde F, Van De Vijver C, Kumar L et al. (2003) Effects of fire and herbivory on the stability of savanna ecosystems. Ecology, 84, 337–350.
- Van Vuuren DP, Edmonds J, Kainuma M et al. (2011) The representative concentration pathways: an overview. Climatic change, 109, 5–31.
- Veldman JW, Putz FE (2011) Grass-dominated vegetation, not species-diverse natural savanna, replaces degraded tropical forests on the southern edge of the Amazon Basin. *Biological Conservation*, 144, 1419–1429.
- Veldman JW, Buisson E, Durigan G et al. (2015a) Toward an old-growth concept for grasslands, savannas, and woodlands. Frontiers in Ecology and the Environment, 13, 154–162.
- Veldman JW, Overbeck GE, Negreiros D et al. (2015b) Where Tree Planting and Forest Expansion are Bad for Biodiversity and Ecosystem Services. *BioScience*, 65, 1011–1018.
- Vodouhê FG, Coulibaly O, Adégbidi A, Sinsin B (2010) Community perception of biodiversity conservation within protected areas in Benin. *Forest Policy and Economics*, 12, 505–512.
- Wessels KJ, Colgan M, Erasmus BFN et al. (2013) Unsustainable fuelwood extraction from South African savannas. Environmental Research Letters, 8, 014007.
- White F (1983) The Vegetation of Africa, a Descriptive Memoir to Accompany the UNESCO/AETFAT/UNSO Vegetation map of Africa. UNESCO, Paris.
- Wigley B, Bond W, Hoffman M (2009) Bush encroachment under three contrasting land-use practices in a mesic South African savanna. *African Journal of Ecology*, 47, 62–70.
- Wittemyer G, Elsen P, Bean WT, Burton ACO, Brashares JS (2008) Accelerated human population growth at protected area edges. *Science*, 321, 123–126.
- Yayneshet T, Treydte A (2015) A meta-analysis of the effects of communal livestock grazing on vegetation and soils in sub-Saharan Africa. *Journal of Arid Environments*, 116, 18–24.
- Zaloumis NP, Bond WJ (2011) Grassland restoration after afforestation: No direction home? *Austral Ecology*, **36**, 357–366.
- Zeng Z, Piao S, Chen A, Lin X, Nan H, Li J, Ciais P (2013) Committed changes in tropical tree cover under the projected 21st century climate change. *Scientific Reports*, **3**, 1–6.
- Zeng Z, Chen A, Piao S, Rabin S, Shen Z (2014) Environmental determinants of tropical forest and savanna distribution: A quantitative model evaluation and its implication. *Journal of Geophysical Research: Biogeosciences*, **119**, 1432– 1445.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Variables included in this study.

Table S2. Changes in forest and savanna area per country for RCPs 2.6 and 4.5 (climate and anthropogenic changes).

Table S3. Changes in forest and savanna areas per country for RCPs 6.0 and 8.5 (climate and anthropogenic changes).

Figure S1. Histograms of tree cover and estimated density of plots.

Figure S2. Partial dependence plots for all the predictors of tree cover using a subset of $N = 10\,000$ points.

Figure S3. Distribution of the WWF Terrestrial Ecoregions (a), protected areas from the World Database on Protected Areas¹ (b), and a classification of the vegetation in sub-Saharan Africa adapted from White's (1983) phytogeographic map (c).

Figure S4. Future projections – RCP 6.0.

Figure S5. Future projections – RCP 8.5.

Figure S6. Difference between current Mean Annual Rainfall (MAR, mm yr⁻¹) from Worldclim 1.4 and MAR estimated from the different RCP scenarios averaged between 2061 and 2080.

Figure S7. Difference between current rainfall seasonality from Worldclim 1.4 and seasonality estimated from the different RCP scenarios averaged between 2061 and 2080.

Figure S8. Difference between current percentage of cropland per pixel and cropland estimated from the different RCP scenarios averaged between 2061 and 2080 [Harmonized Global Land Use version 1 (Chini *et al.*, 2014)].

Figure S9. Difference between current percentage of pasture per pixel and pasture estimated from the different RCP scenarios averaged between 2061 and 2080 [Harmonized Global Land Use version 1 (Chini *et al.*, 2014)].